INDICATIONS AND USAGE

Adjuvant Breast Cancer

ONTRUZANT is indicated for adjuvant treatment of HER2-overexpressing node-positive or nodenegative (ER/PR-negative or with one high-risk feature) breast cancer:

- As part of a treatment regimen containing doxorubicin, cyclophosphamide, and either paclitaxel or docetaxel
- With docetaxel and carboplatin
- As a single agent following multimodality anthracycline-based therapy

Select patients for therapy based on an FDA-approved companion diagnostic for a trastuzumab product.

Metastatic Breast Cancer

ONTRUZANT is indicated:

- In combination with paclitaxel for the first-line treatment of HER2-overexpressing metastatic breast cancer
- As a single agent for treatment of HER2-overexpressing breast cancer in patients who have received one or more chemotherapy regimens for metastatic disease

Select patients for therapy based on an FDA-approved companion diagnostic for a trastuzumab product.

Metastatic Gastric Cancer

ONTRUZANT is indicated, in combination with cisplatin and capecitabine or 5-fluorouracil, for the treatment of patients with HER2 overexpressing metastatic gastric or gastroesophageal junction adenocarcinoma who have not received prior treatment for metastatic disease.

Select patients for therapy based on an FDA-approved companion diagnostic for a trastuzumab product.