

You are Invited!

NINLARO® (ixazomib): An Oral Treatment Option

Presented By

Kasra Karamlou, MD
Assistant Clinical Professor of Medicine
Cleveland Clinic

Hosted by

Takeda

Thursday, May 14, 2020
6:00 PM (ET)

Webcast:

https://event.webcasts.com/starthere.jsp?ei=1311061&tp_key=30e37c5bb6

Sponsored by Takeda. Please note that this is not a continuing medical education (CME) program.

INDICATION

NINLARO® (ixazomib) is indicated in combination with lenalidomide and dexamethasone for the treatment of patients with multiple myeloma who have received at least one prior therapy.

IMPORTANT SAFETY INFORMATION

WARNINGS AND PRECAUTIONS

- **Thrombocytopenia** has been reported with NINLARO. During treatment, monitor platelet counts at least monthly, and consider more frequent monitoring during the first three cycles. Manage thrombocytopenia with dose modifications and platelet transfusions as per standard medical guidelines. Adjust dosing as needed. Platelet nadirs typically occurred between Days 14-21 of each 28-day cycle and recovered to baseline by the start of the next cycle.

Please see additional Important Safety Information continued on page 2 and accompanying NINLARO (ixazomib) full Prescribing Information.

RSVP

via https://event.webcasts.com/starthere.jsp?ei=1311061&tp_key=30e37c5bb6
or by Fax (800) 660-8857
before 05/12/2020

FULL NAME

CREDENTIALS

TITLE

INSTITUTION

CITY

STATE

ZIP CODE

EMAIL ADDRESS

PHONE

LICENSE NUMBER (ONLY IF LICENSED IN MA OR MN)

E20729

MEETING CODE

 **NINLARO**[®]
(ixazomib) capsules
4mg | 3mg | 2.3mg

WARNINGS AND PRECAUTIONS (continued)

- **Gastrointestinal Toxicities**, including diarrhea, constipation, nausea and vomiting, were reported with NINLARO and may occasionally require the use of antidiarrheal and antiemetic medications, and supportive care. Diarrhea resulted in the discontinuation of one or more of the three drugs in 1% of patients in the NINLARO regimen and < 1% of patients in the placebo regimen. Adjust dosing for severe symptoms.
- **Peripheral Neuropathy** (predominantly sensory) was reported with NINLARO. The most commonly reported reaction was peripheral sensory neuropathy (19% and 14% in the NINLARO and placebo regimens, respectively). Peripheral motor neuropathy was not commonly reported in either regimen (< 1%). Peripheral neuropathy resulted in discontinuation of one or more of the three drugs in 1% of patients in both regimens. Monitor patients for symptoms of peripheral neuropathy and adjust dosing as needed.
- **Peripheral Edema** was reported with NINLARO. Monitor for fluid retention. Investigate for underlying causes when appropriate and provide supportive care as necessary. Adjust dosing of dexamethasone per its prescribing information or NINLARO for Grade 3 or 4 symptoms.
- **Cutaneous Reactions**: Rash, most commonly maculo-papular and macular rash, was reported with NINLARO. Rash resulted in discontinuation of one or more of the three drugs in < 1% of patients in both regimens. Manage rash with supportive care or with dose modification.
- **Thrombotic Microangiopathy**: Cases, sometimes fatal, of thrombotic microangiopathy, including thrombotic thrombocytopenic purpura/hemolytic uremic syndrome (TTP/HUS), have been reported in patients who received NINLARO. Monitor for signs and symptoms of TTP/HUS. If the diagnosis is suspected, stop NINLARO and evaluate. If the diagnosis of TTP/HUS is excluded, consider restarting NINLARO. The safety of reinitiating NINLARO therapy in patients previously experiencing TTP/HUS is not known.
- **Hepatotoxicity** has been reported with NINLARO. Drug-induced liver injury, hepatocellular injury, hepatic steatosis, hepatitis cholestatic and hepatotoxicity have each been reported in < 1% of patients treated with NINLARO. Events of liver impairment have been reported (6% in the NINLARO regimen and 5% in the placebo regimen). Monitor hepatic enzymes regularly during treatment and adjust dosing as needed.
- **Embryo-fetal Toxicity**: NINLARO can cause fetal harm. Women should be advised of the potential risk to a fetus, to avoid becoming pregnant, and to use contraception during treatment and for an additional 90 days after the final dose of NINLARO. Women using hormonal contraceptives should also use a barrier method of contraception.

ADVERSE REACTIONS

The most common adverse reactions ($\geq 20\%$) in the NINLARO regimen and greater than the placebo regimen, respectively, were diarrhea (42%, 36%), constipation (34%, 25%), thrombocytopenia (78%, 54%; pooled from adverse events and laboratory data), peripheral neuropathy (28%, 21%), nausea (26%, 21%), peripheral edema (25%, 18%), vomiting (22%, 11%), and back pain (21%, 16%). Serious adverse reactions reported in $\geq 2\%$ of patients included thrombocytopenia (2%) and diarrhea (2%).

DRUG INTERACTIONS: Avoid concomitant administration of NINLARO with strong CYP3A inducers.

SPECIAL POPULATIONS

- **Hepatic Impairment:** Reduce the NINLARO starting dose to 3 mg in patients with moderate or severe hepatic impairment.
- **Renal Impairment:** Reduce the NINLARO starting dose to 3 mg in patients with severe renal impairment or end-stage renal disease requiring dialysis. NINLARO is not dialyzable.
- **Lactation:** Advise nursing women not to breastfeed during treatment with NINLARO and for 90 days after the last dose.

Please see accompanying NINLARO (ixazomib) full Prescribing Information.



All trademarks are the property of their respective owners.

©2020 Millennium Pharmaceuticals, Inc., a wholly owned subsidiary of Takeda Pharmaceutical Company Limited.
All rights reserved. Printed in USA 02/20 MAT-US-IXA-18-00106

