

You are cordially invited to attend a program entitled

PEMAZYRE: The First FDA-Approved Treatment for Adults With Previously Treated, Unresectable, Locally Advanced or Metastatic Cholangiocarcinoma With an FGFR2 Fusion or Other Rearrangement as Detected by an FDA-Approved Test

Tuesday, March 19, 2024 at 6:15 PM

The program will begin at 6:15 PM. Please plan to arrive 15 minutes early to sign in.

Featured Speaker:

Philip Philip, MD

Henry Ford Cancer Institute
Detroit, MI

Location:

Selden Standard

3921 Second Ave
Detroit, MI 48201

Appropriate attendees include licensed HCPs with a direct role in patient care.

Due to a change in Policy, Incyte will no longer provide or pay for alcohol at Speaker Programs.

REGISTRATION

Register by March 12th

<https://sphase.info/inc1141>

To register manually, please contact your Incyte representative Thomas Storti at (810) 280-0486 or tstorti@incyte.com the following information: name, title/degree, state(s) and state license #(s), affiliation, address, phone, and e-mail

Prior to registering, please review the program title and speaker to ensure you have not attended this program within the past year.

Please familiarize yourself with local COVID guidelines prior to attending the program.

INDICATIONS AND USAGE

PEMAZYRE® (pemigatinib) is indicated for the treatment of adults with previously treated, unresectable locally advanced or metastatic cholangiocarcinoma with a fibroblast growth factor receptor 2 (FGFR2) fusion or other rearrangement as detected by an FDA-approved test.

This indication is approved under accelerated approval based on overall response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).

Please see Important Safety Information for PEMAZYRE® on back cover and accompanying Full Prescribing Information.

Please note this program is intended for healthcare professionals (HCPs) only. This program is sponsored by Incyte Corporation and is not eligible for CE credits.

This is an educational event intended only for appropriate healthcare professionals. Spouses, guests, and other individuals who are not the intended audience of this educational program are not permitted to attend. Healthcare professionals who are subject to federal, state or local laws or government ethics restrictions may not attend this event. Incyte will report the cost of any meals provided at this event as required by federal, state or local law requirements.

IMPORTANT SAFETY INFORMATION

Ocular Toxicity

Retinal Pigment Epithelial Detachment (RPED): PEMAZYRE can cause RPED, which may cause symptoms such as blurred vision, visual floaters, or photopsia. Clinical trials of PEMAZYRE did not conduct routine monitoring including optical coherence tomography (OCT) to detect asymptomatic RPED; therefore, the incidence of asymptomatic RPED with PEMAZYRE is unknown.

Among 635 patients who received a starting dose of PEMAZYRE 13.5 mg across clinical trials, RPED occurred in 11% of patients, including Grade 3–4 RPED in 1.3%. The median time to first onset of RPED was 56 days. RPED led to dose interruption of PEMAZYRE in 3.1% of patients, and dose reduction and permanent discontinuation in 1.3% and in 0.2% of patients, respectively. RPED resolved or improved to Grade 1 levels in 76% of patients who required dosage modification of PEMAZYRE for RPED.

Perform a comprehensive ophthalmological examination including OCT prior to initiation of PEMAZYRE and every 2 months for the first 6 months and every 3 months thereafter during treatment. For onset of visual symptoms, refer patients for ophthalmologic evaluation urgently, with follow-up every 3 weeks until resolution or discontinuation of PEMAZYRE. Modify the dose or permanently discontinue PEMAZYRE as recommended in the prescribing information for PEMAZYRE.

Dry Eye: Among 635 patients who received a starting dose of PEMAZYRE 13.5 mg across clinical trials, dry eye occurred in 31% of patients, including Grade 3–4 in 1.6% of patients. Treat patients with ocular demulcents as needed.

Hyperphosphatemia and Soft Tissue Mineralization

PEMAZYRE can cause hyperphosphatemia leading to soft tissue mineralization, cutaneous calcification, calcinosis, and non-uremic calciphylaxis. Increases in phosphate levels are a pharmacodynamic effect of PEMAZYRE. Among 635 patients who received a starting dose of PEMAZYRE 13.5 mg across clinical trials, hyperphosphatemia was reported in 93% of patients based on laboratory values above the upper limit of normal. The median time to onset of hyperphosphatemia was 8 days (range 1–169). Phosphate lowering therapy was required in 33% of patients receiving PEMAZYRE.

Monitor for hyperphosphatemia and initiate a low phosphate diet when serum phosphate level is >5.5 mg/dL. For serum phosphate levels >7 mg/dL, initiate phosphate lowering therapy and withhold, reduce the dose, or permanently discontinue PEMAZYRE based on duration and severity of hyperphosphatemia as recommended in the prescribing information.

Embryo-Fetal Toxicity

Based on findings in an animal study and its mechanism of action, PEMAZYRE can cause fetal harm when administered to a pregnant woman. Oral administration of pemigatinib to pregnant rats during the period of organogenesis caused fetal malformations, fetal growth retardation, and embryo-fetal death at maternal exposures lower than the human exposure based on area under the curve (AUC) at the clinical dose of 13.5 mg.

Advise pregnant women of the potential risk to the fetus. Advise female patients of reproductive potential to use effective contraception during treatment with PEMAZYRE and for 1 week after the last dose. Advise males with female partners of reproductive potential to use effective contraception during treatment with PEMAZYRE and for 1 week after the last dose.

Adverse Reactions: Cholangiocarcinoma

Serious adverse reactions occurred in 45% of patients receiving PEMAZYRE ($n=146$). Serious adverse reactions in $\geq 2\%$ of patients who received PEMAZYRE included abdominal pain, pyrexia, cholangitis, pleural effusion, acute kidney injury, cholangitis infective, failure to thrive, hypercalcemia, hyponatremia, small intestinal obstruction, and urinary tract infection. Fatal adverse reactions occurred in 4.1% of patients, including failure to thrive, bile duct obstruction, cholangitis, sepsis, and pleural effusion.

Permanent discontinuation due to an adverse reaction occurred in 9% of patients who received PEMAZYRE. Adverse reactions requiring permanent discontinuation in $\geq 1\%$ of patients included intestinal obstruction and acute kidney injury.

Dosage interruptions due to an adverse reaction occurred in 43% of patients who received PEMAZYRE. Adverse reactions requiring dosage interruption in $\geq 1\%$ of patients included stomatitis, palmar-plantar erythrodysesthesia syndrome, arthralgia, fatigue, abdominal pain, AST increased, asthenia, pyrexia, ALT increased, cholangitis, small intestinal obstruction, alkaline phosphatase increased, diarrhea, hyperbilirubinemia, electrocardiogram QT prolonged, decreased appetite, dehydration, hypercalcemia, hyperphosphatemia, hypophosphatemia, back pain, pain in extremity, syncope, acute kidney injury, onychomadesis, and hypotension.

Dose reductions due to an adverse reaction occurred in 14% of patients who received PEMAZYRE. Adverse reactions requiring dosage reductions in $\geq 1\%$ of patients who received PEMAZYRE included stomatitis, arthralgia, palmar-plantar erythrodysesthesia syndrome, asthenia, and onychomadesis.

Clinically relevant adverse reactions occurring in $\leq 10\%$ of patients included fractures (2.1%). In all patients treated with pemigatinib, 0.5% experienced pathologic fractures (which included patients with and without cholangiocarcinoma [$N=635$]). Soft tissue mineralization, including cutaneous calcification, calcinosis, and non-uremic calciphylaxis associated with hyperphosphatemia were observed with PEMAZYRE treatment.

Within the first 21-day cycle of PEMAZYRE dosing, serum creatinine increased (mean increase of 0.2 mg/dL) and reached steady state by Day 8, and then decreased during the 7 days off therapy. Consider alternative markers of renal function if persistent elevations in serum creatinine are observed.

In cholangiocarcinoma ($n=146$) the most common adverse reactions (incidence $\geq 20\%$) were hyperphosphatemia (60%), alopecia (49%), diarrhea (47%), nail toxicity (43%), fatigue (42%), dysgeusia (40%), nausea (40%), constipation (35%), stomatitis (35%), dry eye (35%), dry mouth (34%), decreased appetite (33%), vomiting (27%), arthralgia (25%), abdominal pain (23%), hypophosphatemia (23%), back pain (20%), and dry skin (20%).

Drug Interactions

Avoid concomitant use of strong and moderate CYP3A inhibitors with PEMAZYRE. Reduce the dose of PEMAZYRE if concomitant use with a strong or moderate CYP3A inhibitor cannot be avoided. Avoid concomitant use of strong and moderate CYP3A inducers with PEMAZYRE.

Special Populations

Advise lactating women not to breastfeed during treatment with PEMAZYRE and for 1 week after the last dose.

Reduce the recommended dose of PEMAZYRE for patients with severe renal impairment as described in the prescribing information.

Reduce the recommended dose of PEMAZYRE for patients with severe hepatic impairment as described in the prescribing information.

Please see Full Prescribing Information for PEMAZYRE.

PEMAZYRE and the PEMAZYRE logo are registered trademarks of Incyte. Incyte and the Incyte logo are registered trademarks of Incyte. © 2022, Incyte. MAT-PEM-00396 11/22

Pemazyre
(pemigatinib) tablets
13.5mg/9mg/45mg

